CONTENTS

Page

Determination of Acidity and Titrable Nitrogen in Wheat with the Hydrogen Electrode. C. O. SWANSON and E. L. TAGUE.. 1

Ash Absorption by Spinach from Concentrated Soil Solutions. RODNEY H. TRUE, OTIS F. BLACK, and JAMES W. KELLY..... 15

Nitrates, Nitrification, and Bacterial Contents of Five Typical Acid Soils as Affected by Lime, Fertilizer, Crops, and Moisture. H. A. NOYES and S. D. CONNER..... 27

Effect of Certain Ecological Factors on the Morphology of the Urediniospores of Puccinia graminis. E. C. STAKMAN and M. N. LEVINE..... 43

Variations and Mode of Secretion of Milk Solids. JOHN W. GOWEN..... 79

New Biologic Forms of Puccinia graminis. E. C. STAKMAN, M. N. LEVINE, and J. G. LEACH..... 103

Influence of Salts on the Nitric-Nitrogen Accumulation in the Soil. J. E. GREAVES, E. G. CARTER, and H. C. GOLDTHORPE.. 107

Physoderma Disease of Corn. W. H. TISDALE..... 137

Injury to Casuarina Trees in Southern Florida by the Mangrove Borer. THOMAS E. SNYDER..... 155

Life-History Observations on Four Recently Described Parasites of Bruchophagus funebris. THEODORE D. URBAHNS..... 165

Cyanogenesis in Andropogon sorghum. C. T. DOWELL..... 175

Effect of Certain Compounds of Barium and Strontium on the Growth of Plants. J. S. MCHARGUE..... 183

Apple-Scald. CHARLES BROOKS, J. S. COOLEY, and D. F. FISHER. 195

Angular-Leafspot of Tobacco, An Undescribed Bacterial Disease. F. D. FROMME and T. J. MURRAY..... 219

Two Species of Pegomyia Mining the Leaves of Dock. S. W. FROST..... 229

Influence of Foreign Pollen on the Development of Vanilla Fruits. T. B. MCCLELLAND..... 245

A Blood-Destroying Substance in Ascaris lumbricoides. BENJAMIN SCHWARTZ..... 253

Solubility of the Lime, Magnesia, and Potash in Such Minerals as Epidote, Chrysolite, and Muscovite, Especially in Regard to Soil Relationships. R. F. GARDINER..... 259

A Field Study of the Influence of Organic Matter upon the Water-Holding Capacity of a Silt-Loam Soil. FREDERICK J. ALWAY and JOSEPH R. NELLER..... 263

Page

Fusarium-Blight of Potatoes Under Irrigation. H. G. MacMillan 279

Effect of Certain Grain Rations on the Growth of the White Leghorn Chick. G. Davis Buckner, E. H. Nollau, R. H. Wilkins, and Joseph H. Kastle 305

Ammonification of Manure in Soil. H. J. Conn and J. W. Bright 313

Index 351

ERRATA AND AUTHORS' EMENDATIONS

Page 103, line 15, "C I 2879" should read "C I 5879."

Page 104, line 19, "C I 4103" should read "C I 4013."

Page 161, line 31, "*A. labena* n. sp." should read "*Labena* n. sp."

Page 168, the cuts of figures 1 and 2 should be transposed.

Page 229–244, throughout the paper "*Pegomyia affinis* Stein." should read "*Pegomyia vanduse* Mallock 1919."

Page 248, Table III, column 2, line 11, "4¾" should read "3¾."

Page 252, Plate 31, A, should be inverted.

Page 260, Table I, the figures in column 2 should read, from top to bottom, "1.39, 1.94, 1.89, .24, .23, .21. .27, .19, .21, .55"; the figures in column 3 should read ".50, .10, .17, .05, .10, .11, .10, Trace, .03, .10."

Page 261, line 10, "0.27" should read "0.26" and "0.17" should read "0.68."

Page 280, paragraph 2, line 15, "41, B" should read "40, A."

Page 285, paragraph 2, line 5, "38, D" should read "38, C."

Page 286, paragraph 3, line 11, "tube" should read "tuber."

ILLUSTRATIONS

PLATES

NITRATES, NITRIFICATION, AND BACTERIAL CONTENTS OF FIVE TYPICAL ACID SOILS AS AFFECTED BY LIME, FERTILIZER, CROPS, AND MOISTURE

Page

PLATE 1. Representative plates from 1 to 400,000 bacterial dilution of acid yellow silty clay, cropped and held under optimum moisture conditions: A.—Aerobic plates, untreated. B.—Aerobic plates, treated with 2 tons of calcium carbonate. C.—Aerobic plates, treated with complete fertilizer. D.—Aerobic plates, treated with complete fertilizer and 2 tons of calcium carbonate. E.—Aerobic plates, treated with complete fertilizer and 6 tons of calcium carbonate 42

PLATE 2. Representative plates from 1 to 400,000 bacterial dilution of acid whitish silt loam and acid brown silt loam cropped and held under optimum moisture conditions: A.—Aerobic plates, acid whitish silt loam, untreated. B.—Aerobic plates, acid whitish silt loam, treated with 3 tons of calcium carbonate. C.—Aerobic plates, acid whitish silt loam, treated with 500 pounds of acid phosphate. D.—Aerobic plates, acid brown silt loam, untreated. E.—Aerobic plates, acid brown silt loam, treated with 3 tons of calcium carbonate. F.—Aerobic plates, acid brown silt loam, treated with 500 pounds of acid phosphate. G.—Anaerobic plates, acid brown silt loam, untreated. H.—Anaerobic plates, acid brown silt loam, treated with 3 tons of calcium carbonate. I.—Anaerobic plates, acid brown silt loam, treated with 500 pounds of acid phosphate 42

PLATE 3. Representative plates from 1 to 400,000 bacterial dilution of acid black peaty sand, cropped and held under optimum moisture conditions: A.—Aerobic plates, untreated. B.—Aerobic plates, treated with 2 tons of calcium carbonate. C.—Aerobic plates, treated with complete fertilizer. D.—Aerobic plates, treated with complete fertilizer and 2 tons of calcium carbonate. E.—Aerobic plates, treated with complete fertilizer and 6 tons of calcium carbonate. F.—Anaerobic plates, treated with complete fertilizer. G.—Anaerobic plates, treated with complete fertilizer and 2 tons of calcium carbonate. H.—Anaerobic plates, treated with complete fertilizer and 6 tons of calcium carbonate 42

PLATE 4. Representative plates from 1 to 400,000 bacterial dilution of acid dark-brown peat, cropped and held under optimum moisture conditions: A.—Aerobic plates, untreated. B.—Aerobic plates, treated with 2 tons of calcium carbonate. C.—Aerobic plates, treated with 20 tons of calcium carbonate. D.—Anaerobic plates, untreated. E.—Anaerobic plates, treated with 2 tons of calcium carbonate. F.—Anaerobic plates, treated with 20 tons of calcium carbonate 42

PLATE 5. Representative plates from 1 to 40,000 bacterial dilution of acid yellow silty clay kept at different moisture contents: A_2.—Aerobic plates, from soil kept one-half saturated. H_2.—Anaerobic plates, from soil kept one-half saturated. C_2.—Aerobic plates of carbon-dioxid-surviving organisms from soil kept one-half saturated. A_3.—Aerobic plates from soil kept fully saturated. H_3.—Anaerobic plates from soil kept fully saturated. C_3.—Aerobic plates of carbon-dioxid-surviving organisms from soil kept fully saturated 42

Page

PLATE 6. Representative plates from 1 to 40,000 bacterial dilution of acid whitish silt loam kept at different moisture contents: A_1.—Aerobic plates from soil kept one-fourth saturated. H_1.—Anaerobic plates from soil kept one-fourth saturated. C_1.—Aerobic plates of carbon-dioxid-surviving organisms from soil kept one-fourth saturated. A_2.—Aerobic plates from soil kept one-half saturated. H_2.—Anaerobic plates from soil kept one-half saturated. C_2.—Aerobic plates of carbon-dioxid-surviving organisms from soil kept one-half saturated. A_3.—Aerobic plates from soil kept fully saturated. H_3.—Anaerobic plates from soil kept fully saturated. C_3.—Aerobic plates of carbon-dioxid-surviving organisms from soil kept fully saturated. 42

PLATE 7. Representative plates from 1 to 40,000 bacterial dilution of acid brown silt loam kept at different moisture contents: A_1.—Aerobic plates from soil kept one-fourth saturated. H_1.—Anaerobic plates from soil kept one-fourth saturated. C_1.—Aerobic plates of carbon-dioxid-surviving organisms from soil kept one-fourth saturated. A_2.—Aerobic plates from soil kept one-half saturated. H_2.—Anaerobic plates from soil kept one-half saturated. C_2.—Aerobic plates of carbon-dioxid-surviving organisms from soil kept half saturated. A_3.—Aerobic plates from soil kept fully saturated. H_3.—Aerobic plates from soil kept fully saturated. C_3.—Aerobic plates of carbon-dioxid-surviving organisms from soil kept fully saturated. 42

PLATE 8. Representative plates from 1 to 40,000 bacterial dilution of acid black peaty sand kept at different moisture contents: A_1.—Aerobic plates from soil kept one-fourth saturated. H_1.—Anaerobic plates from soil kept one-fourth saturated. C_1.—Aerobic plates of carbon-dioxid-surviving organisms from soil kept one-fourth saturated. A_2.—Aerobic plates from soil kept one-half saturated. H_2.—Anaerobic plates from soil kept one-half saturated. C_2.—Aerobic plates of carbon-dioxid-surviving organisms from soil kept one-half saturated. A_3.—Aerobic plates from soil kept fully saturated. H_3.—Anaerobic plates from soil kept fully saturated. C_3.—Aerobic plates of carbon-dioxid-surviving organisms from soil kept fully saturated. 42

PLATE 9. Representative plates from 1 to 40,000 bacterial dilution of acid dark-brown peat kept at different moisture contents: A_1.—Aerobic plates from soil kept one-fourth saturated. H_1.—Anaerobic plates from soil kept one-fourth saturated. C_1.—Aerobic plates of carbon-dioxid-surviving organisms from soil kept one-fourth saturated. A_2.—Aerobic plates from soil kept one-half saturated. H_2.—Anaerobic plates from soil kept one-half saturated. C_2.—Aerobic plates of carbon-dioxid-surviving organisms from soil kept one-half saturated. A_3.—Aerobic plates from soil kept fully saturated. H_3.—Anaerobic plates from soil kept fully saturated. C_3.—Aerobic plates of carbon-dioxid-surviving organisms from soil kept fully saturated 42

PHYSODERMA DISEASE OF CORN

PLATE A. Corn leaf showing the effects of an attack by *Physoderma zeae-maydis*..... 154

PLATE B. Sheath and culm of corn plant showing the effects of *Physoderma zeae-maydis*..... 154

PLATE 10. Blade and sheath of corn plant showing the effects of severe attack by *Physoderma zeae-maydis*.

Page

PLATE 11. Old sheaths and culms of corn showing effects of severe attacks by *Physoderma zeae-maydis:* A, B.—Badly diseased stalks broken over at weakened, infected lower nodes. C, E.—Portions of old attacked sheaths showing characteristic shredding. D.—Portion of an old infected stalk showing discoloration both on outside and in pith due to the attacks of the fungus 154

PLATE 12. Blades and sheath of corn showing the Physoderma disease produced by inoculating plants in the greenhouse with a suspension of the sporangia of *P. zeae-maydis* in tap water 154

PLATE 13. *Physoderma zeae-maydis:* Various stages in the germination of sporangia, formation of zoospore, and germination of zoospore. *a*.—Sporangium. *b, c, d*.—Opening sporangia showing the early stages of zoospore formation. *e*.—Mature zoospores escaping through the ruptured apex of the endosporangium. *f*.—The collapsing endosporongium after the zoospores have escaped. *g*.—Zoospores. *h*.—Germinating zoospores 154

PLATE 14. *Physoderma zeae-maydis: a-f*.—Zoospores germinating by fine threadlike hyphæ which have penetrated the epidermal cell walls of a tender leaf of Indian corn. In *c, e*, and *f* the enlarged cells have begun to form in the epidermal cells of the host 154

PLATE 15. *Physoderma zeae-maydis:* Mycelial stages within the host cells. *a-d*.—Drawings from ordinary high-power magnifications showing the fibers and enlarged cells of the mycellum. *e-g*.—Drawings magnified with oil-immersion lens. *b, d, g*.—Notice the young sporongia at the ends of the short hyphæ 154

PLATE 16. *Physoderma zeae-maydis: a-e*, Mycelial fibers penetrating the cell walls of the host tissue. *f, g*, Different types of reproductive bodies 154

PLATE 17. *Physoderma zeae-maydis:* Photomicrographs showing the different stages of the development in the host tissue (teosinte). A.—Notice the reproductive bodies connected by the very fine threadlike hyhæ in the central cells of the figure. B.—Host cells filled with mature sporangia ... 154

INJURY TO CASUARINA TREES IN SOUTHERN FLORIDA BY THE MANGROVE BORER

PLATE 18. A.—Casuarina trees planted along the water front, Belle Isle, Miami Beach, Fla., June, 1918. B.—Casuarina trees disfigured and killed by the mangrove borer (*Chrysobathris tranquebarica*) at Miami Beach, Fla. . 164

PLATE 19. *Chrysobothris tranquebarica:* A.—Sex differences in the last abdominal segment. B.—Lateral and dorsal view of ovipositor. C.—Bark of red mangrove (*Rhizophora mangle*) showing how it is divided into plates 164

PLATE 20. *Chrysobothris tranquebarica:* A.—Larval burrow in cambium of Australian pine (*Casurina equisetifolia*), Miami Beach, Fla. B.—Larvae, ventral and dorsal views. C.—Pupa, dorsal and central views. D.—Female and male adult beetles 164

PLATE 21. *Chrysobothris tranquebarica:* Adult male, dorsal view 164

LIFE-HISTORY OBSERVATIONS ON FOUR RECENTLY DESCRIBED PARASITES OF BRUCHOPHAGUS FUNEBRIS

PLATE 22. A.—*Liodontomerus perplexus:* Adult female. B.—*Eutelus bruchophagi:* Adult female 174

PLATE 23. A.—*Trimeromicrus maculatus:* Adult female. B.—*Liodontomerus secundus:* Adult female 174

EFFECT OF CERTAIN COMPOUNDS OF BARIUM AND STRONTIUM ON THE GROWTH OF PLANTS

Page.

PLATE 24. A.—Effect of barium on the growth of cowpeas with and without calcium carbonate. B.—Stimulating effect of barium on root growth of cowpeas. C.—Effect of barium on the growth of soybeans..... 194

ANGULAR-LEAFSPOT OF TOBACCO, AN UNDESCRIBED BACTERIAL DISEASE

PLATE 25. A.—A tobacco leaf showing an early stage of the angular-leafspot. B.—Angular leaf spots on a tobacco leaf..... 228

PLATE 26. A.—Upper surface of a tobacco leaf affected with the angular-leafspot. B.—Lower surface of a tobacco leaf affected with the angular-leafspot..... 228

PLATE 27. A.—Angular leaf spots on a tobacco leaf as seen in transmitted light. B.—Atypical angular leaf spots on a narrow leaf of tobacco..... 228

TWO SPECIES OF PEGOMYIA MINING THE LEAVES OF DOCK

PLATE 28. A.—Eggs of *Pegomyia affinis*. B.—Parasitized eggs of *Pegomyia calyptrata*. C.—Eggs of *Pegomyia calyptrata*. D.—A small mine on Rumex leaf, showing the original linear mine and the beginning of the blotch mine. E.—A typical mine on *Rumex obtusifolius* produced by the larva of *Pegomyia calyptrata*. F.—A monstrosity, an adult *P. calyptrata* issuing feet first from its puparium. G.—A typical mine on *Rumex crispus* produced by the larva of *Pegomyia calyptrata*. H.—A mine on *Rumex obtrusifolius*, produced by a nearly mature larva entering a new leaf to complete its development..... 244

PLATE 29. A.—Fiske cages, used in studying the adult flies of *Pegomyia* spp., showing the arrangement in outdoor insectary. B.—Glass cylinder cage, used in studying habits of adult flies. C.—Fiske cage, used in studying habits of adult flies..... 244

PLATE 30. A.—Ventral aspect of posterior segment of larva of *Pegomyia affinis*, showing lobes and anal opening. B.—Lateral aspect of posterior segment of larva of *P. affinis*, showing tubercles. C.—Lateral aspect of posterior segment of larva of *P. calyptrata*. D.—Ventral aspect of posterior segment of larva of *P. calyptrata*, showing lobes and anal opening. E.—"Pseudocephalon" (Henneguy) of *P. calyptrata*: *a*, buttonlike areas referred to in text; *b*, sensory papilla; *c*, antenna; *d*, mandibular sclerite; *e*, anterior spiracle. F.—Posterior spiracle of larva of *P. calyptrata*..... 244

INFLUENCE OF FOREIGN POLLEN ON THE DEVELOPMENT OF VANILLA FRUITS

PLATE 31. A.—Left to right, *Vanilla planifolia*, first of each pair close-fertilized, second fertilized with vanillon pollen. B.—Cross sections of same fruits..... 252

PATE 32. A.—Vanillon No. 13 fruits, close- and cross-fertilized. Lower row close-fertilized, upper fertilized with pollen of *Vanilla planifolia*. B.—Cross section of same fruits..... 252

PLATE 33. A.—Vanillon No. 34: Fruits at right fertilized with pollen of *Vanilla planifolia*, left close-fertilized. B.—Cross sections of same fruits. 252

PLATE 34. A.—Vanillon No. 43 fruits. Upper row fertilized with pollen of *Vanilla planifolia*, lower row close-fertilized. B.—Cross sections of same fruits..... 252

Page.

PLATE 35. A.—Longitudinal sections of vanilla fruits. Left to right: Vanillon, pistillate, × vanillon, staminate; vanillon, pistillate, × *Vanilla planifolia*, staminate; *Vanilla planifolia*, pistillate, × vanillon, staminate; *Vanilla planifolia*, pistillate, × *Vanilla planifolia*, staminate. B.—Comparative length of cleared columns and ovaries. Vanillon above, *Vanilla planifolia* below..... 252

A FIELD STUDY OF THE INFLUENCE OF ORGANIC MATTER UPON THE WATER-HOLDING CAPACITY OF A SILT-LOAM SOIL

PLATE 36. View of Field J., Minn. Agr. Experiment Station Farm, showing topography and surroundings, looking from plot 6 to the barn which occupies part of plot 1..... 278

FUSARIUM-BLIGHT OF POTATOES UNDER IRRIGATION

PLATE 37. Effect of Fusarium-blight on seed pieces of potato: A.—Early Ohio seed pieces: Control above; pieces inoculated with *F. oxysporum* below. B.—Early Ohio plant. See piece inoculated with *F. oxysporum*. C.—Early Ohio seed pieces: Control (left) and inoculated (right) seed pieces. D.—Seed piece well decayed, resulting from soil infection. E.—Seed-piece rot in field..... 304

PLATE 38. A.—Inoculated and uninoculated stems of same potato plant. B.—Potato plant (control), showing method of inoculating with wedge of melilotus stem. C.—Seed-piece rot in field. D.—Potato plant naturally infected by *F. oxysporum* in the field..... 304

PLATE 39. Potato stems showing seed-piece rot: A.—Stem split to show rotting due to organism entering through seed piece from soil. B.—Stem split to show slight discoloration at base where infection from soil-infected seed piece occurred. C.—Seed piece of potato plant shown in Plate 40, B. D.—Seed-piece rot in field..... 304

PLATE 40. Potato plants affected by Fusarium blight: A.—Potato plant late in season with rolled leaves. Fusarium blight. B.—Top of potato plant consisting of three stems. C.—Potato plant of two stems; one at left showing Fusarium blight, or wilt caused by seed-piece infection; one at right healthy. D.—Plant with rolled leaves gradually dying from Fusarium blight..... 304

PLATE 41. A.—A field of potatoes showing the result of unfavorable cultural and soil conditions, by which seed-piece rot destroyed 60 per cent of the stand. B.—A field of potatoes planted with whole seed (rows to right) and cut seed (rows to left) at midseason..... 304

EFFECT OF CERTAIN GRAIN RATIONS ON THE GROWTH OF THE WHITE LEGHORN CHICK

PLATE 42. A.—Lot 1 at the age of 3 months. B.—Lot 3 at the age of 3 months. C.—Lot 2 at the age of 3 months..... 312

TEXT FIGURES

DETERMINATION OF ACIDITY AND TITRABLE NITROGEN IN WHEAT WITH THE HYDROGEN ELECTRODE

FIG. 1. Graphs showing the hydrogen-ion concentration of wheat extract on titration to P_H 7..... 6

2. Graphs showing the hydrogen-ion concentration of wheat extract on titration P_H 8.3..... 6

106547°—20——2

Page

FIG. 3. Graphs showing the hydrogen-ion concentration of wheat extract on titration to P_H 9.3..... 6
4. Graphs showing the production of amino nitrogen at different temperatures..... 10
5. Graphs showing the total phosphorus in wheat extract at different periods of extraction and at different temperatures..... 12
6. Graphs showing the inorganic phosphorus in wheat extract at different periods of extraction and at different temperatures..... 12

ASH ABSORPTION BY SPINACH FROM CONCENTRATED SOIL SOLUTIONS

FIG. 1. Graphs representing total ash and individual ash constituents found in spinach tops from plots receiving the substances indicated..... 20
2. Graphs representing total ash and individual ash constituents in spinach roots from plots receiving the substances indicated..... 20

NITRATES, NITRIFICATION, AND BACTERIAL CONTENTS OF FIVE TYPICAL ACID SOILS AFFECTED BY LIME, FERTILIZER, CROPS, AND MOISTURE

FIG. 1. Graphs showing the relation of aerobes and anaerobes to nitrification of five acid soils with and without lime and fertilizer treatments..... 37
2. Graphs showing the relation of aerobes and anaerobes to nitrates in five acid soils kept at different moisture contents..... 38

INFLUENCE OF SALTS ON THE NITRIC-NITROGEN ACCUMULATION IN THE SOIL

FIG. 1. Graphs showing molecular concentrations at which the highest stimulation is noted..... 128
2. Graphs showing the percentage of stimulation at the above noted molecular concentrations (see fig. 1), the untreated soil being counted as producing 100 per cent of nitric nitrogen..... 129
3. Graphs showing the molecular concentrations at which the various salts are toxic to nitrification..... 130
4. Graphs showing the molecular concentrations which reduce the nitrification to three-fourths normal..... 131
5. Graphs showing the percentages of nitric nitrogen produced in 100 gm. of soil to which had been added 2×10^{-3} mole of the various salts, the untreated soil being counted as producing 100 per cent..... 132

PHYSODERMA DISEASE OF CORN

FIG. 1. Map showing the distribution of *Physoderma zeae-maydis* in the United States. Broken lines, *P. zeae-maydis* present: solid line, *P. zeae-maydis* causing damage..... 138

INJURY TO CASUARINA TREES IN SOUTHERN FLORIDA BY THE MANGROVE BORER

FIG. 1. *Chrysobothris tranquebarica:* Larva, dorsal, lateral, and ventral views... 159
2. *Chrysobothris tranquebarica: a*, Female pupa, ventral view; *b*, same, dorsal view..... 160

LIFE-HISTORY OBSERVATIONS ON FOUR RECENTLY DESCRIBED PARASITES OF BRUCHOPHAGUS FUNEBRIS

Page

FIG. 1. *Liodontomerus perplexus:* Larva 168
2. *Liodontomerus perplexus:* Pupa 168
3. *Liodontomerus secundus:* Larva 170
4. *Liodontomerus secundus:* Pupa 170
5. *Eutelus bruchophagi:* Larva 171
6. *Eutelus bruchophagi:* Pupa 171
7. *Trimeromicrus maculatus:* Larva 173
8. *Trimeromicrus maculatus:* Pupa 173

APPLE-SCALD

FIG. 1. Graphs showing the effect of maturity upon susceptibility of Grimes apples to scald 197
2. Graphs showing the relative susceptibility to scald of eastern and western Grimes apples 198
3. Graphs showing the effects of temperature on apple-scald at the end of 4, 6, 7, 9, 13, and 16 weeks 199
4. Graphs showing the effects of temperature on apple-scald at the end of 2, 3, 4, 5, 7, 12, and 16 weeks 200
5. Graphs showing the effects of temperature on apple-scald at the end of 2, 3, 4, 5, 9, and 11 weeks 202
6. Graphs showing the effect of temperature on apple-scald at the end of 5, 7, 9, 11, 13, 16, and 19 weeks 203
7. Graphs showing the effect of temperature on apple-scald at the end of 7, 10, 12, and 18 weeks 204
8. Graphs showing the effect of temperature on apple-scald at the end of 12, 14, 20, and 26 weeks 205
9. Graphs showing the effect of temperature on apple-scald at the end of 8, 12, 16, and 20 weeks 205
10. Graphs showing the effect of temperature on apple-scald at the end of 8, 12, and 16 weeks 209
11. Graphs showing the relation of carbon dioxid to apple-scald production. 210

TWO SPECIES OF PEGOMYIA MINING THE LEAVES OF DOCK

FIG. 1. *A*, pharyngeal skeleton of first instar, *Pegomyia calpytrata* Zett; *B*, pharyngeal skeleton of second instar, *P. calyptrata;* C, mandibular sclerite of second instar, *P. calyptrata*; *D*, pharyngeal skeleton of third instar, *P. calyptrata; E*, pharyngeal skeleton of first instar, *P. affinis* Stein; *F*, mandibular sclerite of first instar, *P. affinis; G*, pharyngeal skeleton of second instar, *P. affinis; H*, pharyngeal skeleton of third instar, *P. affinis* 233

A FIELD STUDY OF THE INFLUENCE OF ORGANIC MATTER UPON THE WATER-HOLDING CAPACITY OF A SILT-LOAM SOIL

FIG. 1. Diagram showing arrangement of plots and crops on field J, University Farm, St. Paul, Minn. *A* is the original plan adhered to from 1893 to 1914, while *B* shows the cropping plan in 1915. *C* shows the arrangement of the samples taken for the two composites from a plot 265
2. Diagram showing the amount and distribution of the rainfall at University Farm, St. Paul, Minn., during part of the season of 1915. The dates of sampling are indicated by asterisks 269